



Clinical Practice Guidelines: Trauma/Traumatic haemorrhagic shock

Policy code	CPG_TR_THS_0924
Date	September, 2024
Purpose	To ensure a consistent approach to the management of a patient with a traumatic brain injury.
Scope	Applies to Queensland Ambulance Service (QAS) clinical staff.
Health care setting	Pre-hospital assessment and treatment.
Population	Applies to all ages unless stated otherwise.
Source of funding	Internal – 100%
Author	Clinical Quality & Patient Safety Unit, QAS
Review date	September, 2027
Information security	UNCLASSIFIED – Queensland Government Information Security Classification Framework.
URL	https://ambulance.qld.gov.au/clinical.html

While the QAS has attempted to contact all copyright owners, this has not always been possible. The QAS would welcome notification from any copyright holder who has been omitted or incorrectly acknowledged.

All feedback and suggestions are welcome. Please forward to: Clinical.Guidelines@ambulance.qld.gov.au

Disclaimer

The Digital Clinical Practice Manual is expressly intended for use by appropriately qualified QAS clinicians when performing duties and delivering ambulance services for, and on behalf of, the QAS.

The QAS disclaims, to the maximum extent permitted by law, all responsibility and all liability (including without limitation, liability in negligence) for all expenses, losses, damages and costs incurred for any reason associated with the use of this manual, including the materials within or referred to throughout this document being in any way inaccurate, out of context, incomplete or unavailable.

© State of Queensland (Queensland Ambulance Service) 2024.



This work is licensed under the **Creative Commons Attribution-NonCommercial-NoDerivatives V4.0 International License**

You are free to copy and communicate the work in its current form for non-commercial purposes, as long as you attribute the State of Queensland, Queensland Ambulance Service and comply with the licence terms. If you alter the work, you may not share or distribute the modified work. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>

For copyright permissions beyond the scope of this license please contact: Clinical.Guidelines@ambulance.qld.gov.au

Traumatic haemorrhagic shock

September, 2024

Haemorrhage is the primary cause of preventable mortality in people with traumatic injuries. It is estimated to be responsible for 40% of all deaths related to trauma.^[1] Substantial ongoing bleeding remains a critical and potentially fatal complication of severe trauma.

Initial treatment of these patients must prioritise rapid haemorrhage control achieved through the application of direct pressure bandages, tourniquets, wound packing, splinting, and/or pelvic binders.^[2] This is followed by the maintenance of normothermia in addition to avoidance of both acidosis and coagulopathy.^[3,4] Additionally pharmacological adjuncts and blood products that aim to inhibit fibrinolysis, replace clotting factors, and restore circulatory volume should then be considered.

The circulating blood volume varies between individuals depending on their weight and gender. In clinical practice, the following estimated can be used:

- Adult male: 75 mL/kg
- Adult female: 65 mL/kg
- Paediatric: 80 mL/kg

Clinical features



- The clinical features of haemorrhagic shock vary depending on the amount of blood loss.

SIGNS AND SYMPTOMS OF HAEMORRHAGE				
Parameter	Minor	Mild	Moderate	Severe
Approximate blood loss	< 15%	15–30%	31–40%	> 40%
Heart rate	↔	↔/↑	↑	↑/↑↑
Blood pressure	↔	↔	↔/↓	↓
Pulse pressure	↔	↓	↓	↓
Respiratory rate	↔	↔	↔/↑	↑
Urine output	↔	↔	↓	↓↓
Glasgow Coma Scale score	↔	↔	↓	↓
Need for blood products	Monitor	Possible	Yes	Massive Transfusion Protocol

* Base excess is the quantity of base (HCO_2^- , in mEq/L) that is above or below the normal range in the body. A negative number is called a base deficit and indicates metabolic acidosis.

LEGEND:

↔	↑	↓	↑↑	↓↓
no change	increase	decrease	significant increase	significant decrease



Risk assessment

- Blood loss in the out-of-hospital setting is inherently difficult to gauge and is often underestimated.
- Hypotension in trauma patients may not be secondary to haemorrhage – consider other causes (e.g., obstructive shock, tension pneumothorax, cardiac tamponade, spinal cord injuries, or toxins).



Additional information

- Prior to definitive care, the out-of-hospital fluid resuscitation goal for most patients with traumatic haemorrhagic shock is the maintenance of a palpable radial pulse. Alternatively, in patients suffering an associated traumatic brain injury and a potential for raised intracranial pressure, the target becomes supporting a systolic blood pressure of 100–120 mmHg.
- It's well-recognised that a significant cohort of traumatically injured patients develop clotting dysfunction. Whilst the exact pathophysiological mechanism of trauma-induced coagulopathy (TIC) is not fully understood,^[6] it is likely to be influenced by genetic factors, injury burden, hypoperfusion, and resuscitative measures.^[7]
- Hypofibrinogenemia is one of the earliest manifestations of coagulation dysfunction and is associated with increased mortality and an increased requirement of blood transfusions.^[4,8]
- Low fibrinogen is associated with reduced clot strength and poor outcomes in trauma patients.^[9] Hypofibrinogenemia in the prehospital setting and on emergency department arrival has been associated with both increased mortality and increased blood transfusions.^[4,10]
- Identifying trauma patients who require blood products and have low plasma fibrinogen concentrations allows for the administration of human fibrinogen substitution therapy.

CPG: Clinician safety
CPG: Standard cares

Note: Clinicians must only perform procedures for which they have received specific training and authorisation by the QAS.

- **Haemorrhage control**
- **Maintain normothermia**

Consider:

- **Oxygen**
- **IV/IO access**
- **Sodium chloride 0.9%**
- **Tranexamic acid**
- **PRBC (1 unit to 2.2 mmol calcium gluconate)**
- **ELP (1 unit to 2.2 mmol calcium gluconate)**
- **Human fibrinogen**

Associated traumatic brain injury?

N

Treatment target
= palpable radial pulse

Y

Treatment target
= SBP 100–120 mmHg

Transport to hospital
Pre-notify as appropriate